

Cooling of Neutron Stars with Color Superconducting Quark Cores

Hovik Grigorian^{1,2}, David Blaschke^{3,4}, and Dmitri Voskresensky^{5,6}

¹*Fachbereich Physik, Universität Rostock, D-18051 Rostock, Germany*

²*Department of Physics, Yerevan State University,*

Alex Manoogian Str. 1, 375025 Yerevan, Armenia

³*Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany*

⁴*Bogoliubov Laboratory of Theoretical Physics,*

Joint Institute for Nuclear Research, 141980, Dubna, Russia

⁵*Theory Division, GSI mbH, D-64291 Darmstadt, Germany*

⁶*Moscow Institute for Physics and Engineering, 115409 Moscow, Russia*

Abstract

We show that within a recently developed nonlocal chiral quark model the critical density for a phase transition to color superconducting quark matter under neutron star conditions can be low enough for these phases to occur in compact star configurations with masses below $1.3 M_{\odot}$. We study the cooling of these objects in isolation for different values of the gravitational mass. Our equation of state (EoS) allows for 2SC quark matter with a large quark gap ~ 100 MeV for u and d quarks of two colors that coexists with normal quark matter within a mixed phase in the hybrid star interior. We argue that, if the phases with unpaired quarks were allowed, the corresponding hybrid stars would cool too fast. If they occurred for $M < 1.3 M_{\odot}$, as it follows from our EoS, one could not appropriately describe the neutron star cooling data existing by today. We discuss a "2SC+X" phase, as a possibility to have all quarks paired in two-flavor quark matter under neutron star constraints, where the X-gap is of the order of 10 keV - 1 MeV. Density independent gaps do not allow to fit the cooling data. Only the presence of an X-gap that decreases with increase of the density could allow to appropriately fit the data in a similar compact star mass interval to that following from a purely hadronic model. This scenario is suggested as an alternative explanation of the cooling data in the framework of a hybrid star model.

I. INTRODUCTION

Recently, we have reinvestigated the cooling of neutron stars within a purely hadronic model [1], i.e., suppressing the possibility of quark cores in neutron star interiors. We have demonstrated that the neutron star cooling data available by today can be well explained within the "*Nuclear medium cooling scenario*", cf. [2, 3], i.e., if one includes medium effects into consideration. This scenario puts a tight constraint to the density dependence of the asymmetry energy in the nuclear equation of state (EoS). The latter dependence is an important issue for the analysis of heavy ion collisions especially within the new CBM (compressed baryon matter) program to be realized at the future accelerator facility FAIR at GSI Darmstadt. The density dependence of the asymmetry energy determines the proton fraction in neutron star matter and thus governs the onset of the very efficient direct Urca (DU) process. This process, once occurring, would lead to a very fast cooling of neutron stars. Thereby, if neutron stars with $M < M_{\text{crit}}^{\text{DU}}$ cooled slowly, the stars with the mass M only slightly above $M_{\text{crit}}^{\text{DU}}$ would be already characterized by a very rapid cooling, yielding surface temperatures significantly below those given by the X -ray data points. Unfortunately, the mass spectrum of cooling neutron stars is not known. One can not rely on neutron star mass measurements in binary radio pulsars (BRPs) with the averaged neutron star mass value $M_{\text{BRP}} = 1.35 \pm 0.04 M_{\odot}$ [4, 5] because these objects come from a twice selected population: probably not all neutron stars go through the active radio-pulsar stage [6], and the properties of neutron stars in binaries may be different from those of isolated objects, see e.g. [7]. However, it seems not a fair assumption that most of cooling neutron stars should be essentially lighter than the lower limit for BRP masses $M_{\text{BRP,min}} = 1.31 M_{\odot}$. Therefore, the threshold for the DU process should better occur at high enough densities such that typical neutron stars are not affected by it. This assumption about the mass distribution can be developed into a more quantitative test of cooling scenarios when these are combined with population synthesis models. The latter allow to obtain Log N – Log S distributions for nearby coolers which can be tested with data from the ROSAT catalogue [8]. In addition the microscopically based Argonne model for the EoS, that is supposed to be the most realistic one [9], yields a critical mass for the DU process $\sim 2 M_{\odot}$. At these high star masses the central baryon density exceeds already the 5-fold of the nuclear saturation density such that exotic states of matter as, e.g., hyperonic matter or quark matter are more

appropriate. Ref. [10] argued that the presence of quark matter in massive compact star cores is a most reliable hypothesis since within their model hyperons appear at sufficiently low densities and the equation of state becomes too soft to carry neutron star masses in the range of M_{BRP} [11].

For quark matter in neutron stars, the interaction ($\alpha_s \neq 0$) permits the quark DU process. The works [12, 13, 14, 15] have shown that this result does not exclude the possibility that neutron stars might possess large quark matter cores. The suppression of the emissivity of the DU process may arise due to the occurrence of diquark pairing and thus color superconductivity in low-temperature quark matter. To demonstrate this possibility one has used the simplifying assumption that all quarks are gapped with the same gap which is not valid for superconducting phases at low densities. In the present paper we abandon that restriction and suggest a more subtle pairing pattern that is constrained by comparison with neutron star cooling phenomenology.

II. COLOR SUPERCONDUCTIVITY

The quark-quark interaction in the color anti-triplet channel is attractive driving the pairing with a large zero-temperature pairing gap $\Delta \sim 100$ MeV for the quark chemical potential $\mu_q \sim (300 \div 500)$ MeV, cf. [16, 17], for a review see [18] and references therein. The attraction comes either from the one-gluon exchange, or from a non-perturbative 4-point interaction motivated by instantons [19], or from non-perturbative gluon propagators [20]. Various phases are possible. The so called 2-flavor color superconductivity (2SC) phase allows for unpaired quarks of one color, say blue. There may also exist a color-flavor locked (CFL) phase [21] for not too large values of the dynamical strange quark mass or in other words for very large values of the baryon chemical potential [22], where the color superconductivity is complete in the sense that the diquark condensation produces a gap for quarks of all three colors and flavors. The value of the gap is of the same order of magnitude as that in the two-flavor case. There exist other attractive quark pairing channels. However the attraction is rather weak. One of these channels allows for a color spin locking (CSL), where all quarks are paired in a spin-1 channel. The weak pairing channels are characterized by gaps typically in the interval $\sim 10 \text{ keV} \div 1 \text{ MeV}$.

III. HYBRID STARS

In describing the hadronic part of the hybrid star, as in [1], we exploit the Argonne $V18 + \delta v + UIX^*$ model of the EoS given in [9], which is based on the most recent models for the nucleon-nucleon interaction with the inclusion of a parameterized three-body force and relativistic boost corrections. Actually we continue to adopt an analytic parameterization of this model by Heiselberg and Hjorth-Jensen [23], hereafter HHJ, that smoothly incorporates causality at high densities. The HHJ EoS fits the symmetry energy to the original Argonne $V18 + \delta v + UIX^*$ model in the mentioned density interval yielding the threshold density for the DU process $n_c^{\text{DU}} \simeq 5.19 n_0$ ($M_c^{\text{DU}} \simeq 1.839 M_\odot$).

The 2SC phase occurs at lower baryon densities than the CFL phase, see [24, 25]. For applications to compact stars the omission of the strange quark flavor is justified by the fact that chemical potentials in central parts of the stars do barely reach the threshold value at which the mass gap for strange quarks breaks down and they appear in the system [26].

We will employ the EoS of a nonlocal chiral quark model developed in [27] for the case of neutron star constraints with a 2SC phase. It has been shown in that work that the Gaussian formfactor ansatz leads to an early onset of the deconfinement transition and such a model is therefore suitable to discuss hybrid stars with large quark matter cores [28].

In some density interval above the onset of the first order phase transition, there may appear a mixed phase region, see [29]. Ref. [29] disregarded finite size effects, such as surface tension and charge screening. Refs [30] on the example of the hadron-quark mixed phase have demonstrated that finite size effects might play a crucial role substantially narrowing the region of the mixed phase or even forbidding its appearance in a hybrid star configuration. Therefore we omit the possibility of the hadron-quark mixed phase in our model assuming that the quark phase arises by the Maxwell construction. In this case we found a tiny density jump on the boundary from $n_c^{\text{hadr}} \simeq 0.44 \text{ fm}^{-3}$ to $n_c^{\text{quark}} \simeq 0.46 \text{ fm}^{-3}$.

A large difference between chemical potentials of u and d quarks forbids the pure 2SC phase, cf. [27]. The CFL phase is still not permitted at such densities. Ignoring weak pairing patterns there could be two possibilities: either the quark matter is in the normal phase, or there appears a region of the 2SC – normal quark mixed phase. Disregarding finite size effects, as in [29], Refs. [25, 28] have argued for a broad region of the 2SC – normal quark mixed phase instead of a pure 2SC phase. In our model the mixed phase continues

up to the central density n_{centr} . Thus we have the mixed phase in the density interval: $n_c^{hadr} < n_c^{(mix,1)} < n_c^{quark} < n < n_c^{(mix,2)} = n_{centr}$. In the given case the arguments of [30] might be partially relaxed since the surface tension for the 2SC – normal quark boundary should be significantly smaller than for the quark – hadron boundary. Indeed, as has been indicated in Ref. [15], the surface tension is proportional to the difference of the energies of two phases, being proportional to $(\Delta/\mu_q)^2 \ll 1$ in the given case. Recently, Ref. [31] estimated the surface tension to be typically smaller than 5 MeV/fm² for densities of our interest. For the choice of the Gaussian formfactor the mixed phase appears for sufficiently low density $n > n_c^{(mix,1)} \simeq n_c^{hadr} = 0.44 \text{ fm}^{-3}$ ($M > M_c^{quark} = 1.214 M_\odot$). The presence or absence of the 2SC - normal quark mixed phase instead of only one of those phases is not so important for the hybrid star cooling problem since the latter is governed by processes involving either normal excitations or excitations with the smallest gap.

In Fig. 1 we present the mass-radius relation for hybrid stars with HHJ EoS vs. Gaussian nonlocal chiral quark separable model (SM) EoS. Configurations with the 2SC - normal quark mixed phase, “HHJ-SM with 2SC” (HHJ hadronic EoS together with Gaussian nonlocal chiral quark model SM permitting 2SC - normal quark mixed phase) given by the solid line, are stable, whereas without 2SC large gap color superconducting matter (“HHJ-SM without 2SC”) no stable hybrid star configuration is possible. In the case “HHJ-SM with 2SC” the maximum neutron star mass proves to be $1.793 M_\odot$. For an illustration of the constraints on the mas-radius relation which can be derived from compact star observations we show in Fig. 1 the compactness limits from the thermal emission of the isolated neutron star RX J1856.5-3754 as given in [32] and from the redshifted absorption lines in the X-ray burst spectra of EXO 0748-676 given in [33]. These are, however, rather weak constraints.

Additionally, within the “HHJ-SM with 2SC” mixed phase we will allow for the possibility of a weak pairing channel for all the quarks which were unpaired, with typical gaps $\Delta_X \sim 10 \text{ keV} \div 1 \text{ MeV}$, as in the case of the CSL pairing channel, see [34, 35]. Since we don’t know yet the exact pairing pattern for this case, we call this hypothetical phase “2SC+X”. In such a way all the quarks get paired, some strongly in the 2SC channel and some weakly in the X channel.

IV. COOLING

For the calculation of the cooling of the hadron part of the hybrid star we use the same model as in [1]. The main processes are the medium modified Urca (MMU) and the pair breaking and formation (PBF) processes. The HHJ EoS is adopted. Below we choose two cooling models used in [1] distinguished by different sets of the nucleon gaps. The first choice (further model I) is shown in Fig. 2 (Fig. 12 of Ref. [1]). We use a fit for the relation between surface and crust temperatures (called “our fit” in [1]). We also made calculations for the other two crust models used in Ref. [1]. Although the results are affected by the choice of the crust model, our conclusions remain untouched. The possibilities of pion condensation and of other so called exotic processes are suppressed since in the model [1] these processes may occur only for neutron star masses exceeding $M_c^{\text{quark}} = 1.214 M_\odot$. The DU process is irrelevant in this model up to very large neutron star mass $M > 1.839 M_\odot$. The $1S_0$ neutron and proton gaps are taken the same as those shown by thick lines in Fig. 5 of Ref. [1]. We pay particular attention to the fact that the $3P_2$ neutron gap is additionally suppressed by the factor 0.1 compared to that shown in Fig. 5 of [1]. The latter suppression is motivated by the result of the recent work [36] and is required to fit the cooling data.

For the calculation of the cooling of the quark core in the hybrid star we use the model [14]. We incorporate the most efficient processes: the quark direct Urca (QDU) processes on unpaired quarks, the quark modified Urca (QMU), the quark bremsstrahlung (QB), the electron bremsstrahlung (EB), and the massive gluon-photon decay (see [12]). Following [37] we include the emissivity of the quark pair formation and breaking (QPFB) processes. The specific heat incorporates the quark contribution, the electron contribution and the massless and massive gluon-photon contributions. The heat conductivity contains quark, electron and gluon terms.

We are basing on the picture presented in Fig. 2 and add the contribution of the quark core. For the Gaussian formfactor the quark core occurs already for $M > 1.214 M_\odot$ according to the model [27], see Fig. 1. Most of the relevant neutron star configurations (see Fig. 2) are then affected by the presence of the quark core. First we check the possibility of the 2SC+ normal quark phases.

Fig. 3 shows the cooling curves calculated with the given Gaussian ansatz. The variation of the gaps for the strong pairing of quarks within the 2SC phase and the gluon-photon

mass in the interval $\Delta, m_{g-\gamma} \sim 20 \div 200$ MeV only slightly affects the results. The main cooling process is the QDU process on normal quarks. We see that the presence of normal quarks entails too fast cooling. The data could be explained only if all the masses lie in a very narrow interval ($1.21 < M/M_\odot < 1.22$ in our case). In case of the other two crust models the resulting picture is similar. The existence of only a very narrow mass interval in which the data can be fitted seems us unrealistic as by itself as from the point of view of the observation of the neutron stars in binary systems with different masses, e.g., $M_{\text{B1913+16}} \simeq 1.4408 \pm 0.0003 M_\odot$ and $M_{\text{J0737-3039B}} \simeq 1.250 \pm 0.005 M_\odot$, cf. [5]. *Thus the data can't be satisfactorily explained.*

In Fig. 4 we permit the weak pairing pattern for all quarks which before in Fig. 3 were assumed to be unpaired. We use $\Delta_X \simeq 1$ MeV for the corresponding quark gap. Fig. 4 demonstrates too slow cooling. Even slow cooling data are not explained. The same statement holds for all three crust models. Thus the gaps for formerly unpaired quarks should be still smaller in order to obtain a satisfactory description of the cooling data.

In Fig. 5 we again allow for the weak pairing pattern of those quarks which were assumed to be unpaired in Fig. 3, but now we use $\Delta_X = 30$ keV for the corresponding residual quark pairing gap (X). We see that all data points, except the Vela, CTA 1 and Geminga, correspond to hybrid stars with masses in the narrow interval $M = 1.21 \div 1.22 M_\odot$, very similar to that in Fig. 3. This also seems to be unrealistic with the same argument as for the presence of unpaired quarks, see Fig. 3 above. We have obtained a similar picture for the other two crust models. We have varied the constant Δ in wide limits but the qualitative picture does not change.

Therefore we would like to explore whether a density-dependent X-gap could allow a description of the cooling data within a larger interval of compact star masses. We don't choose the X-gap as a rising function of the chemical potential since the feature of such a model will not spread the accessible mass interval satisfying the cooling data constraint relative to the density independent X-gap case. We employ the ansatz

$$\Delta_X(\mu) = \Delta_c \exp[-\alpha(\mu - \mu_c)/\mu_c] , \quad (1)$$

where the parameters are chosen such that at the critical quark chemical potential $\mu_c = 330$ MeV for the onset of the deconfinement phase transition the X-gap has its maximal value of $\Delta_c = 1.0$ MeV and at the highest attainable chemical potential $\mu_{\text{max}} = 507$ MeV, i.e. in

the center of the maximum mass hybrid star configuration it falls to a value of the order of 10 keV. We choose the value $\alpha = 10$ for which $\Delta_X(\mu_{\max}) = 4.6$ keV. In Fig. 6 we show the resulting cooling curves for the ansatz (1). We observe that the mass interval for compact stars which obey the cooling data constraint ranges now from $M = 1.32 M_\odot$ for slow coolers up to $M = 1.75 M_\odot$ for fast coolers such as Vela, cf. with that we have found with the purely hadronic model [1] with different parameter choices. Note that according to a recently suggested independent test of cooling models [8] by comparing results of a corresponding population synthesis model with the Log N - Log S distribution of nearby isolated X-ray sources the cooling model I did not pass the test. Thereby it would be interesting to see whether our quark model within the gap ansatz (1) may pass the Log N - Log S test.

In [1] different hadronic cooling models have been presented which can explain the available cooling data by varying the neutron star mass in some limits. The question arises whether one of these alternatives could be preferred. In Ref. [1] we conjectured that the model based on Fig. 20 of [1] (Fig. 7 of the given work) that has used another choice of the gaps demonstrates the best fit since it shows a more regular dependence on the neutron star masses. We call it model II. In this model II the $1S_0$ neutron gap is the same as in the model I used above, whereas the proton $1S_0$ gap is calculated by [38] (see thin lines in Fig. 5 of [1]). The neutron $3P_2$ gap is taken from [38] but then is additionally suppressed by factor 0.1 as required by the result of [36] and in order to appropriately fit the cooling data. According to [8] this model II successfully passed the mentioned Log N - Log S test supporting the conjecture of [8] and an earlier conjecture of [39] that neutron star masses should be essentially different.

We have performed the same hybrid cooling calculation as described above, now for the model II. The results are qualitatively the same as those presented in Figs 3 – 6. The difference is that now the curves related to low mass objects ($M < M_{\text{crit}}^{\text{DU}}$) are pushed downwards describing the Crab data point. However, as in model I, both scenarios either that there are unpaired quarks and that the X-gaps are density independent, do not allow to appropriately fit the data.

The results for the density-dependent X-gap according to Eq. (1) now using the hadronic cooling model II (demonstrated by Fig. 7) are shown in Fig. 8. In comparison with the result of Fig. 6 the variation of the cooling data between slow and fast coolers is obtained by choosing hybrid star masses in a wider range $1.1 \leq M/M_\odot \leq 1.7$. Next, it would be

interesting to check whether this model passes the Log N - Log S test.

V. CONCLUSION

Concluding, we demonstrated that within a recently developed nonlocal, chiral quark model the critical densities for a phase transition to color superconducting quark matter can be low enough for these phases to occur in compact star configurations with masses below $1.3 M_{\odot}$. For the choice of the Gaussian formfactor the 2SC - normal quark matter mixed phase arises at $M \simeq 1.21 M_{\odot}$. We have shown that without a residual pairing the 2SC quark matter phase could describe the cooling data only if compact stars had masses in a very narrow band around the critical mass for which the quark core can occur. Since there are observations of neutron stars with higher and essentially different masses such a scenario should be disfavored.

Then we assumed that formally unpaired quarks can be paired with small gaps $\Delta_X < 1$ MeV (2SC+X pairing), which values we varied in wide limits. With density independent gaps we failed to appropriately fit the cooling data.

We showed that the present day cooling data could be still explained by hybrid stars, however, when assuming a complex pairing pattern, where quarks are partly strongly paired within the 2SC channel, and partly weakly paired with gaps $\Delta_X < 1$ MeV, being rapidly decreasing with the increase of the density. Within the given hypothesis and in the framefork of our quark model for the equation of state the observational data are explained in a broad mass interval $M = 1.32 - 1.75 M_{\odot}$ for the model I and $M = 1.1 - 1.7 M_{\odot}$ for the model II, as it is favored by the experimental data. However we will point out that the result is model dependent and demonstrates only a possibility that the presence of the 2SC+X phase might not contradict to the neutron star cooling data.

We have found rather strong constraints for the density dependent X-gap pairing. It remains to be investigated which microscopic pairing pattern could fulfill the constraints obtained in this work. Another indirect check of the model could be the Log N - Log S test.

Acknowledgement

The authors acknowledge the hospitality and support of Rostock University where most of this work has been performed. The work of H.G. has been supported in part by the Virtual Institute of the Helmholtz Association under grant No. VH-VI-041 and by DAAD University partnership program; that of D.V. has been supported in part by DFG grant No. 436 RUS 17/117/03.

-
- [1] D. Blaschke, H. Grigorian, and D.N. Voskresensky, Cooling of Neutron Stars. Hadronic Model, *Astron. Astrophys.* **424**, 979 (2004).
 - [2] C. Schaab, D. Voskresensky, A.D. Sedrakian, F. Weber, and M.K. Weigel, *Astron. Astrophys.* **321**, 591 (1997).
 - [3] D.N. Voskresensky, in book: "Physics of Neutron Star Interiors", *Lecture Notes in Physics*, Eds. D. Blaschke, N.K. Glendenning, A. Sedrakian, Springer, Heidelberg (2001), p. 467-502.
 - [4] S. E. Thorsett, and D. Chakrabarty, *Astrophys. J.* **512**, 288 (1999).
 - [5] I.H. Stairs, *Science* **304**, 547 (2004).
 - [6] E. V. Gotthelf, and G. Vasisht, in *Proc. IAU Coll. 177 "Pulsar Astronomy - 2000 and beyond"*, eds. M. Kramer, N. Wex, and N. Wielebinski, *ASP Conf. Series* 202, 699 (2001).
 - [7] P. Podsiadlowski, et al., *Astrophys. J.* **612**, 1044 (2004).
 - [8] S. Popov, H. Grigorian, R. Turolla, and D. Blaschke, *arXiv:astro-ph/0411618*.
 - [9] A. Akmal, V.R. Pandharipande, and D.G. Ravenhall, *Phys. Rev. C* **58**, 1804 (1998).
 - [10] M. Baldo, G. F. Burgio and H. J. Schulze, *arXiv:astro-ph/0312446*.
 - [11] This statement essentially depends on the poorly known nucleon-hyperon interaction.
 - [12] D. Blaschke, T. Klähn, and D.N. Voskresensky, *Astrophys. J.* **533**, 406 (2000).
 - [13] D. Page, M. Prakash, J.M. Lattimer and A. Steiner, *Phys. Rev. Lett.* **85** 2048 (2000).
 - [14] D. Blaschke, H. Grigorian and D.N. Voskresensky, *Astron. & Astrophys.* **368**, 561 (2001).
 - [15] D. Blaschke, D.N. Voskresensky and H. Grigorian, *arXiv:astro-ph/0403171*.
 - [16] M. Alford, K. Rajagopal, and F. Wilczek, *Phys. Lett. B* **422**, 247 (1998).
 - [17] R. Rapp, T. Schäfer, E.V. Shuryak, and M. Velkovsky, *Phys. Rev. Lett.* **81**, 53 (1998).
 - [18] K. Rajagopal, and F. Wilczek, "The Condensed Matter Physics of QCD", in: B.L. Ioffe

- Festschrift *At the Frontier of Particle Physics / Handbook of QCD*, vol. 3, edited by M. Shifman, World Scientific, Singapore, 2001, p. 2061.
- [19] D. Diakonov, H. Forkel, and M. Lutz, Phys. Lett. **B 373**, 147 (1996); G.W. Carter, and D. Diakonov, Phys. Rev. **D 60**, 016004 (1999); R. Rapp, E. Shuryak, and I. Zahed, Phys. Rev. **D 63**, 034008 (2001).
 - [20] D. Blaschke, and C.D. Roberts, Nucl. Phys. **A 642**, 197 (1998); J.C.R. Bloch, C.D. Roberts, and S.M. Schmidt, Phys. Rev. **C 60**, 65208 (1999).
 - [21] M. Alford, K. Rajagopal, and F. Wilczek, Nucl. Phys. **B 357**, 443 (1999); T. Schäfer, and F. Wilczek, Phys. Rev. Lett. **82**, 3956 (1999).
 - [22] M. Alford, J. Berges, and K. Rajagopal, Nucl. Phys. **B 558**, 219 (1999).
 - [23] H. Heiselberg, and M. Hjorth-Jensen, Astrophys. J. **525**, L45 (1999).
 - [24] A.W. Steiner, S. Reddy, and M. Prakash, Phys. Rev. **D 66**, 094007 (2002).
 - [25] F. Neumann, M. Buballa, and M. Oertel, Nucl. Phys. **A714**, 481 (2003).
 - [26] C. Gocke, D. Blaschke, A. Khalatyan, and H. Grigorian, arXiv:hep-ph/0104183.
 - [27] D. Blaschke, S. Fredriksson, H. Grigorian, and A. Öztas, Nucl. Phys. **A 736**, 203 (2004).
 - [28] H. Grigorian, D. Blaschke, and D.N. Aguilera, Phys. Rev. **C 69**, 065802 (2004).
 - [29] N.K. Glendenning, Phys. Rev. **D 46** 1274 (1992); N.K. Glendenning, Phys. Rep. **342**, 393 (2001).
 - [30] D.N. Voskresensky, M. Yasuhira, and T. Tatsumi, Phys. Lett. **B 541**, 93 (2002); Nucl. Phys. **A 723**, 291 (2002).
 - [31] S. Reddy, and G. Rupak, arXiv:nucl-th/0405054.
 - [32] M. Prakash, J.M. Lattimer, A.W. Steiner, and D. Page, Nucl. Phys. **A 715**, 835 (2003).
 - [33] J. Cottam, F. Paerels, and M. Mendez, Nature **420**, 51 (2002).
 - [34] T. Schäfer, Phys. Rev. D **62**, 094007 (2000).
 - [35] A. Schmitt, Q. Wang and D. H. Rischke, Phys. Rev. D **66**, 114010 (2002).
 - [36] A. Schwenk, and B. Friman, Phys. Rev. Lett. **92**, 082501 (2004).
 - [37] P. Jaikumar, and M. Prakash, Phys. Lett. **B 516**, 345 (2001).
 - [38] T. Takatsuka, and R. Tamagaki, Prog. Theor. Phys. **112**, 37 (2004).
 - [39] D.N. Voskresensky, and A.V. Senatorov, JETP. **63**, 885 (1986).




FIG. 1: Mass - radius relations for compact star configurations with different EoS: purely hadronic star with HHJ EoS (dashed line), stable hybrid stars with HHJ - Gaussian nonlocal chiral quark separable model (SM) with 2SC phase (solid line) and with HHJ - SM, without 2SC phase (dash-dotted line). Data for two sources are also indicated, see text.




FIG. 2: Model I. Cooling curves according to the nuclear medium cooling scenario, see Fig. 12 of [1]. The labels correspond to the gravitational masses of the configurations in units of the solar mass.




FIG. 3: Model I. Cooling curves for hybrid star configurations with Gaussian quark matter core in the 2SC phase. The labels correspond to the gravitational masses of the configurations in units of the solar mass.




FIG. 4: Model I. Cooling curves for hybrid star configurations with Gaussian quark matter core in the 2SC+X phase. The weak pairing gap is $\Delta_X \simeq 1$ MeV. The labels correspond to the gravitational masses of the configurations in units of the solar mass.




FIG. 5: Same as Fig. 4 but with a weak pairing gap of 30 keV.




FIG. 6: Same as Fig. 4 but with a density dependent pairing gap according to Eq. (1).




FIG. 7: Model II. Same as Fig. 20 of Ref. [1].




FIG. 8: Same as Fig. 6 but for model II.